

PROTOCOLLO EFFETTI LUNGO TERMINE

1. Porta D, Milani S, Lazzarino AI, Perucci CA, Forastiere F. Systematic review of epidemiological studies on health effects associated with management of solid waste. *Environ Health*. 2009 Dec 23;8:60. doi: 10.1186/1476-069X-8-60.Review.
2. Risultati del progetto Moniter: (http://www.arpa.emr.it/pubblicazioni/moniter/generale_1526.asp)
3. Gorla S, C. Daniau, et al. (2009). "Risk of cancer in the vicinity of municipal solid waste incinerators: importance of using a flexible modelling strategy." *Int J Health Geogr* 8: 31.
4. Federico M, M. Pirani, et al. (2010). "Cancer incidence in people with residential exposure to a municipal waste incinerator: an ecological study in Modena (Italy), 1991-2005." *Waste Manag* 30(7): 1362-1370
5. Ranzi A, Fano V, Erspamer L, Lauriola P, Perucci CA, Forastiere F. Mortality and morbidity among people living close to incinerators: a cohort study based on dispersion modeling for exposure assessment. *Environ Health*. 2011 Mar 24;10:22. doi: 10.1186/1476-069X-10-22.
6. Gouveia N, Prado RR. Spatial analysis of the health risks associated with solid waste incineration: a preliminary analysis. *Rev Bras Epidemiol*. 2010 Mar;13(1):3-10.
7. Viel JF, Daniau C, Gorla S, Fabre P, de Crouy-Chanel P, Sauleau EA, Empereur-Bissonnet P. Risk for non Hodgkin's lymphoma in the vicinity of French municipal solid waste incinerators. *Environ Health*. 2008;7:51.
8. Viel JF, Arveux P, Baverel J, Cahn JY. Soft-tissue sarcoma and non-Hodgkin's lymphoma clusters around a municipal solid waste incinerator with high dioxin emission levels. *Am J Epidemiol*. 2000;152:13-19.
9. Floret N, Mauny F, Challier B, Arveux P, Cahn JY, Viel JF. Dioxin emissions from a solid waste incinerator and risk of non-Hodgkin lymphoma. *Epidemiology*. 2003;14:392-398.
10. Comba P, Ascoli V, Belli S, Benedetti M, Gatti L, Ricci P, Tieghi A. Risk of soft tissue sarcomas and residence in the neighborhood of an incinerator of industrial wastes. *Occup Environ Med*. 2003;60:650-683.
11. Zambon P, Ricci P, Bovo E, Casula A, Gattolin M, Fiore AR, Chiuosi F, Guzzinati S. Sarcoma risk and dioxin emissions from incinerators and industrial plants: a population-based case-control study (Italy). *Environ Health*. 2007;6:19.
12. Floret N, Mauny F, Challier B, Cahn JY, Tourneux F, Viel JF. Dioxin emissions and soft-tissue sarcoma: results of a population-based case-control study. *Rev Epidemiol Sante Publique*. 2004 Jun;52(3):213-20.
13. Elliott P, Shaddick G, Kleinschmidt I, Jolley D, Walls P, Beresford J, Grundy C. Cancer incidence near municipal solid waste incinerators in Great Britain. *Br J Cancer*. 1996;73:702-710
14. Viel JF, Clement MC, Hägi M, Grandjean S, Challier B, Danzon A. Dioxin emissions from a municipal solid waste incinerator and risk of invasive breast cancer: a population-based case-control study with GIS-derived exposure. *Environ Health*. 2008;7:4. doi: 10.1186/1476-069X-7-51
15. Elliott P, Eaton N, Shaddick G, Carter R. Cancer incidence near municipal solid waste incinerators in Great Britain. Part 2: histopathological and case-note review of primary liver cancer cases. *Br J Cancer*. 2000;82:1103-1106.
16. Biggeri A, Barbone F, Lagazio C, Bovenzi M, Stanta G. Air pollution and lung cancer in Trieste, Italy: spatial analysis of risk as a function of distance from sources. *Environ Health Perspect*. 1996;104:750-754.
17. Parodi S, Baldi R, Benco C, Franchini M, Garrone E, Vercelli M, Pensa F, Puntoni R, Fontana V. Lung cancer mortality in a district of La Spezia (Italy) exposed to air pollution from industrial plants. *Tumori*. 2004;90:181-185.
18. Knox E. Childhood cancers, birthplaces, incinerators and landfill sites. *Int J Epidemiol*. 2000;29:391-397.
19. Dummer TJ, Dickinson HO, Parker L. Adverse pregnancy outcomes around incinerators and crematoriums in Cumbria, north west England, 1956-93. *J Epidemiol Community Health*. 2003;57:456-461.
20. Cordier S, Chevrier C, Robert-Gnansia E, Lorente C, Brula P, Hours M. Risk of congenital anomalies in the vicinity of municipal solid waste incinerators. *Occup Environ Med*. 2004;61:8-15.
21. Cordier S, A. Lehebel, et al. (2010). "Maternal residence near municipal waste incinerators and the risk of urinary tract birth defects." *Occup Environ Med* 67(7): 493-499.

22. Jansson B, Voog L. Dioxin from Swedish municipal incinerators and the occurrence of cleft lip and palate malformations. *Int J Environ Stud.* 1989;34:99-104.
23. ten Tusscher, G. W., G. A. Stam, et al. (2000). "Open chemical combustions resulting in a local increased incidence of orofacial clefts." *Chemosphere* 40(9-11): 1263-1270.
24. Tango T, Fujita T, Tanihata T, Minowa M, Doi Y, Kato N, Kunikane S, Uchiyama I, Tanaka M, Uehata T. Risk of adverse reproductive outcomes associated with proximity to municipal solid waste incinerators with high dioxin emission levels in Japan. *J Epidemiol.* 2004;14:83-93.
25. Cresswell, P. A., J. E. S. Scott, et al. (2003). "Risk of congenital anomalies near the Byker waste combustion plant." *Journal of Public Health Medicine* 25(3): 237-242.
26. Vinceti, M., C. Malagoli, et al. (2008). "Adverse pregnancy outcomes in a population exposed to the emissions of a municipal waste incinerator." *Science of the Total Environment* 407(1): 116-121.
27. Lloyd OL, Lloyd MM, Williams FL, Lawson A. Twinning in human populations and in cattle exposed to air pollution from incinerators. *Br J Ind Med.* 1988;45:556-560.
28. Williams FL, Lawson AB, Lloyd OL. Low sex ratios of births in areas at risk from air pollution from incinerators, as shown by geographical analysis and 3- dimensional mapping. *Int J Epidemiol.* 1992;21:311-319.
29. Lin C-H, Lin C-Y, et al. (2006). "Birth outcomes of infants born in areas with elevated ambient exposure to incinerators generated PCDD/F. *Environ Int* 32: 624-629".
30. Hsiue TR, Lee SS, Chen HI. Effects of air pollution resulting from wire reclamation incineration on pulmonary function in children. *Chest.* 1991;100:698-702.
31. Shy CM, Degnan D, Fox DL, Mukerjee S, Hazucha MJ, Boehlecke BA, Rothenbacher D, Briggs PM, Devlin RB, Wallace DD, Stevens RK, Bromberg PA. Do waste incinerators induce adverse respiratory effects? An air quality and epidemiological study of six communities. *Environ Health Perspect.* 1995;103:714-724.
32. Lee JT, Shy CM. Respiratory function as measured by peak expiratory flow rate and PM10: six communities study. *J Expo Anal Environ Epidemiol.* 1999;9:293- 299.
33. Miyake Y, Yura A, Misaki H, Ikeda Y, Usui T, Iki M, Shimizu T. Relationship between distance of schools from the nearest municipal waste incineration plant and child health in Japan. *Eur J Epidemiol.* 2005;20:1023-1029.
34. Franchini M, Rial M, Buiatti E, Bianchi F. Health effects of exposure to waste incinerator emissions:a review of epidemiological studies. *Ann Ist Super Sanita.* 2004;40(1):101-15.
35. Minichilli F, Bartolacci S, Buiatti E, Bianchi F. Evaluation of the health impact of a waste-to-energy incinerator. *Epidemiol Prev.* 2007 Jan-Feb;31(1 Suppl 2):35-41.
36. Linzalone N, Bianchi F. Incinerators: not only dioxins and heavy metals, also fine and ultrafine particles. *Epidemiol Prev.* 2007 Jan-Feb;31(1):62-6.
37. Forastiere F, Perucci CA. Wastes, scientific evidence and policy making: which role for epidemiology?. *Epidemiol Prev.* 2008 Jan-Feb;32(1):5-7.
38. Pirastu R, Comba P. SENTIERI Project analyzed mortality of 5.5 million Italians in polluted sites. *Epidemiol Prev.* 2011 Sep-Dec;35(5-6 Suppl 2):108-9.