

Mortality in Children and Adolescents, 1990–2013

Top 5 Causes of Death by Age Group in Developed and Developing Countries, 2013

DEVELOPED COUNTRIES


RANK	AGE <1 y	AGE 1–4 y	AGE 5–9 y	AGE 10–19 y
1	Congenital anomalies	Congenital anomalies	Road injuries	Road injuries
2	Preterm birth complications	Road injuries	Drowning	Self-harm
3	Other neonatal disorders	Drowning	Congenital anomalies	Interpersonal violence
4	Neonatal encephalopathy	LRI	Brain cancer	Drowning
5	Sudden infant death syndrome	Fire and heat	Leukemia	Other neoplasms

DEVELOPING COUNTRIES

RANK	AGE <1 y	AGE 1–4 y	AGE 5–9 y	AGE 10–19 y
1	Preterm birth complications	Malaria	Diarrhea	Road injuries
2	LRI	LRI	LRI	HIV/AIDS
3	Neonatal encephalopathy	Diarrhea	Intestinal infectious diseases	Self-harm
4	Congenital anomalies	Protein-energy malnutrition	Malaria	Drowning
5	Neonatal sepsis	Drowning	Road injuries	Intestinal infectious diseases

LRI=lower respiratory tract infections

Death Rates by Age Group by World Health Organization (WHO) Region, 1990–2013



WHO region definitions: http://www.who.int/healthinfo/global_burden_disease/definition_regions/en/

Authors: Laurie Marczak, PhD; Kevin O'Rourke, MFA; Dawn Shepard, BA, for the Institute for Health Metrics and Evaluation.

Sources: GBD 2013 Pediatrics Collaboration. *JAMA Pediatr.* 2016; 170(3):267-287.
doi: 10.1001/jamapediatrics.2015.4276
Please cite as: *JAMA.* 2016;315(19):2055.
10.1001/jama.2016.5891

